

MATH 210: Construction of the Reals Homework

1. Let C^∞ be the set of all continuous functions on the interval $[0, 1]$. Show that $d(f, g) = \int_0^1 |f(x) - g(x)| dx$ is a metric on C^∞ .
2. Give two other elements (as Cauchy sequences) of the equivalence class $\{1, \frac{1}{10}, \frac{1}{100}, \frac{1}{1000} \dots\} / =_C$.
3. Give the element of \mathbb{R} (in standard notation) that corresponds to

$$\left\{ 1, 1 + \frac{1}{1}, 1 + \frac{1}{1 + \frac{1}{1}}, 1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{1}}}, \dots \right\} / =_C$$

4. A sequence $\{x_n\}$ is said to be **eventually bounded by M** iff there exists N such that for every $n > N$ we have $|x_n| < M$. Prove that if $\{x_n\}$ is a Cauchy sequence then it is eventually bounded.
(Hint: Since the sequence is Cauchy, it has a limit, call it L . Then use the limit definition of $x_n \rightarrow L$ to show that there exists N such that for $n > N$ we have $L - 1 < x_n < L + 1$. Now find the appropriate bound on $|x_n|$.)
5. Suppose $\{r_n\} =_C \{s_n\}$ and $\{t_n\} =_C \{u_n\}$. Show that
 - (a) $\{r_n + t_n\} =_C \{s_n + u_n\}$
 - (b) $\{r_n t_n\} =_C \{s_n u_n\}$ (Hint: use $0 = -s_n t_n + s_n t_n$ and Problem 4.)

6. Consider the sequence of rationals $\{1, 4, 16, \dots, 4^n, \dots\}$ and the metric on \mathbb{Q} given by

$$d_2(x, y) = 2^{(\# \text{ of } 2\text{'s in the denominator of } x-y) - (\# \text{ of } 2\text{'s in the numerator of } x-y)}.$$

- (a) Prove that this sequence is Cauchy under the metric $d_2(x, y)$.
- (b) Show that this sequence has limit $L = 0$ under the metric $d_2(x, y)$.